




Fluid
therapy.org

Chapter 34:

Gastrointestinal Losses and Upper GI Bleeding



To get a copy of the book, visit: www.fluidtherapy.org

34

Gastrointestinal Losses and Upper GI Bleeding

Composition of Gastrointestinal Secretions 403

VOMITING

Fluid and Electrolyte Abnormalities.. 404

Management..... 406

Etiological treatment..... 406

Correction of hypovolemia and electrolyte disturbances..... 406

Fluid selection for gastric irrigation..... 407

DIARRHEA

Fluid and Electrolyte Abnormalities.. 408

Management..... 410

Etiological treatment..... 410

Correction of hypovolemia and electrolyte disturbances..... 410

Oral rehydration therapy.....411

Intravenous fluid therapy411

UPPER GASTROINTESTINAL BLEEDING

Common Presentations 412

Management..... 413

Initial assessment..... 413

Hemodynamic assessment 413

Basic laboratory investigations 413

History and evaluation..... 413

Treatment 413

Stabilization..... 413

Blood transfusion..... 414

Medications 415

Placement of nasogastric tube 417

Endoscopy..... 417

Specific treatment..... 417

Risk factors..... 417

Monitoring..... 418

Loss of gastrointestinal (GI) secretions and hematemesis are frequently encountered problems in clinical practice. Several gastrointestinal disorders lead to loss of GI secretions causing electrolytes and acid-base disturbances. The composition of gastrointestinal secretions and potential disturbances due to their losses are summarized in Table 34.1 [1–3], which will help plan its management.

The most common disorders, vom-

iting, diarrhea, and hematemesis, will be discussed.

VOMITING

Vomiting or nasogastric suction is a common problem in clinical practice causing loss of fluid and electrolytes. Therefore, it is necessary to understand the basic pathophysiology of hypovolemia and electrolytes and acid-base abnormali-

ties due to upper GI losses and the proper selection of IV fluids to correct it.

FLUID AND ELECTROLYTE ABNORMALITIES

It is important to understand the fluid and electrolyte abnormalities that commonly occur due to vomiting.

Gastric juice contains approximately: Sodium (Na^+) 20–60 mEq/L, chloride (Cl^-) 140 mEq/L, potassium (K^+) 14

mEq/L, and hydrogen ions (H^+) 60–80 mEq/L [2].

So, vomiting (or nasogastric-NG suction) leads to hypokalemic, hypochloremic metabolic alkalosis with hypovolemia. Due to the loss of HCl containing gastric juice, the first abnormalities usually occur in vomiting are hypochloremia with metabolic alkalosis, and hypokalemia occurs subsequently. The pathogenesis of these abnormalities is summarized below (Figure 34.1).

Want to read more?

Get Printed Version

Get Kindle Version

REFERENCES

1. Gennari FJ, Weise WJ. Acid-base disturbances in gastrointestinal disease. *Clin J Am Soc Nephrol*. 2008;3(6):1861–8.
2. National Clinical Guideline Centre (UK). *Intravenous Fluid Therapy: Intravenous Fluid Therapy in Adults in Hospital* [Internet]. London: Royal College of Physicians (UK); 2013.
3. Do C, Evans GJ, DeAguero J, et al. Dysnatremia in Gastrointestinal Disorders. *Front. Med*. 2022;9:892265.
4. National Collaborating Centre for Women's and Children's Health (UK). *Diarrhoea and Vomiting Caused by Gastroenteritis: Diagnosis, Assessment and Management in Children Younger than 5 Years*. London: RCOG Press; 2009. (NICE Clinical Guidelines, No. 84.) 5, Fluid management. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK63837/>.
5. Shah GS, Das BK, Kumar S, et al. Acid base and electrolyte disturbance in diarrhoea. *Kathmandu Univ Med J (KUMJ)*. 2007;5(1):60–62.
6. Onyiriuka AN, Iheagwara EC. Serum electrolyte profiles of under-five Nigerian children admitted for severe dehydration due to acute diarrhea. *Niger J Heal Sci* 2015;15(1):14–17.
7. Dhyani A, Ameta P, Patel JB, et al. Clinical profile of children with diarrhoea admitted in pediatric intensive care unit of Bal Chikitsalay, M.B. Hospital, RNT Medical College, Udaipur, Rajasthan, India. *Int J Contemp Pediatr* 2016;3(4):1371–1374.
8. Dastidar RG, Konar N. A Study of Electrolyte Disturbances in a Child Presenting with Acute Gastroenteritis, with Special Emphasis on Hyponatremic Dehydration-A Hospital based CrossSectional Study. *Pediatr Ther* 2017;7(2):322.
9. Gopchade A. Electrolyte Disturbances in Children Presenting with Acute Diarrhea: A Prospective Cohort Study. *JMSCR* 2019;7(9):777.
10. Tayab MA, Hoq MA. Acid-base and Electrolyte Disturbances in Children Presenting with Acute Watery Diarrhoea in Emergency Observation and Referral Unit of Dhaka Shishu (Children) Hospital. *Dhaka Shishu (Children) Hospital Journal*, 2021;36(2):120–124.
11. Riaz L, Hussain MK, Javed M, et al. Challenges and Complications Associated with Sodium, Potassium Imbalances and Preventive Measures in Children Under-Five with Acute Gastroenteritis. *Pak Armed Forces Med J* 2022;72(Suppl-1):S64–67.
12. Kiela PR, Ghishan FK. Physiology of Intestinal Absorption and Secretion. *Best Pract Res Clin Gastroenterol*. 2016;30(2):145–159.

13. Nagami GT. Hyperchloremia - Why and How. *Nefrologia*. 2016;36(4):347–53.
14. Mahajan V, Sajan SS, Sharma A, et al. Ringer's lactate vs normal saline for children with acute diarrhea and severe dehydration: A double blind randomized controlled trial. *Indian Pediatr* 2012;49(1):963–68.
15. Naseem M, Dubey AP, Mishra TK, et al. Effect of Rehydration with Normal Saline versus Ringer Lactate on Serum Sodium Level of Children with Acute Diarrhea and Severe Dehydration: A Randomized Controlled Trial. *Indian Pediatr*. 2020;57(6):519–522.
16. World Health Organization. The treatment of diarrhoea: a manual for physicians and other senior health workers, 4th rev. World Health Organization. 2005. <https://apps.who.int/iris/handle/10665/43209>.
17. Mahalanabis D, Brayton JB, Mondal A, et al. The use of Ringer's lactate in the treatment of children with cholera and acute noncholera diarrhoea. *Bull World Health Organ*. 1972;46(3):311–9.
18. Cieza JA, Hinostroza J, Huapaya JA, et al. Sodium chloride 0.9% versus Lactated Ringer in the management of severely dehydrated patients with choleraform diarrhoea. *J Infect Dev Ctries*. 2013;7(7):528–32.
19. Rasheed S, Rafique S, Hussain A, et al. Comparison of outcome between Ringer's lactate and normal saline fluid replacement in pediatric patients with acute watery diarrhoea. *PAFMJ*. 2020;70(Suppl-1):S101–05.
20. Shaikh B, Ghoto IS, Langah AU, et al. A Randomized Double Blind Controlled Trial of Ringers Lactate versus Normal Saline among Pediatric Patients with Acute Severe Diarrhea. *PJMHS* 2022;16(3):281–282.
21. Lobo DN, Awad S. Should chloride-rich crystalloids remain the mainstay of fluid resuscitation to prevent “pre-renal” acute kidney injury? *con*. *Kidney International*. 2014;86(6):1096–1105.
22. Heng Li, Sun SR, Yap JQ, et al. 0.9% saline is neither normal nor physiological. *J Zhejiang Univ Sci B*. 2016;17(3):181–187.
23. Yunus NM, Bellomo R, Glassford N, et al. Chloride-liberal vs. Chloride-restrictive intravenous fluid administration and acute kidney injury: an extended analysis. *Intensive Care Med* 2015;41:257–64.
24. Kamboj AK, Hoversten P, Leggett CL. Upper Gastrointestinal Bleeding: Etiologies and Management. *Mayo Clin Proc*. 2019;94(4):697–703.
25. Cañamares-Orbis P, Lanás Arbeloa Á. New Trends and Advances in Non-Variceal Gastrointestinal Bleeding-Series II. *J Clin Med*. 2021;10(14):3045.
26. Laine L, Barkun AN, Saltzman JR, et al. ACG Clinical Guideline: Upper Gastrointestinal and Ulcer Bleeding. *Am J Gastroenterol*. 2021;116(5):899–917.
27. Tielleman T, Bujanda D, Cryer B. Epidemiology and risk factors for upper gastrointestinal bleeding. *Gastrointest Endosc Clin N Am*. 2015;25(3):415–428.
28. Wuerth BA, Rockey DC. Changing Epidemiology of Upper Gastrointestinal Hemorrhage in the Last Decade: A Nationwide Analysis. *Dig Dis Sci*. 2018;63(5):1286–1293.
29. Oakland K. Changing epidemiology and etiology of upper and lower gastrointestinal bleeding. *Best Pract Res Clin Gastroenterol* 2019;42–43:101610.
30. Wilkins T, Wheeler B, Carpenter M. Upper Gastrointestinal Bleeding in Adults: Evaluation and Management. *Am Fam Physician*. 2020 Mar 1;101(5):294–300. Erratum in: *Am Fam Physician*. 2021;103(2):70.
31. Cappell MS, Friedel D. Initial management of acute upper gastrointestinal bleeding: from initial evaluation up to gastrointestinal endoscopy. *Med Clin North Am* 2008;92(3):491–509.
32. Kupfer Y, Cappell MS, Tessler S. Acute gastrointestinal bleeding in the intensive care unit. The intensivist's perspective. *Gastroenterol Clin North Am* 2000;29(2):275–307.
33. Elsayed IAS, Battu PK, Irving S. Management of acute upper GI bleeding. *BJA Education* 2017;17(4):117–123.
34. Dinesen L, Benson M. Managing acute upper gastrointestinal bleeding in the acute assessment unit. *Clin Med (Lond)*. 2012;12(6):589–593.
35. Hernandez-Gea V, Berbel C, Baiges A, et al. Acute variceal bleeding: risk stratification and management (including TIPS). *Hepatol Int* 2018;12:81–90.
36. Villanueva C, Colomo A, Bosch A. Transfusion for acute upper gastrointestinal bleeding. *N Engl J Med*. 2013;368(14):1362–3.
37. Jairath V, Kahan BC, Gray A, et al. Restrictive versus liberal blood transfusion for acute upper gastrointestinal bleeding (TRIGGER): A pragmatic, open-label, cluster randomised feasibility trial. *Lancet* 2015;386(9989):137–44.
38. Carson JL, Stanworth SJ, Roubinian N, et al. Transfusion thresholds and other strategies for guiding allogeneic red blood cell transfusion. *Cochrane Database Syst Rev* 2016;10:CD002042.
39. Odutayo A, Desborough MJR, Trivella M et al. Restrictive versus liberal blood transfusion for gastrointestinal bleeding: a systematic review and meta-analysis of randomised controlled trials. *Lancet Gastro-enterol Hepatol* 2017;2(5):354–360.
40. Carson JL, Stanworth SJ, Dennis JA, et al. Transfusion thresholds for guiding red blood cell transfusion. *Cochrane Database Syst Rev* 2021;12(12):CD002042.
41. Carson JL, Guyatt G, Heddle NM, et al. Clinical practice guidelines from the AABB: Red blood cell transfusion thresholds and storage. *JAMA* 2016;316(19):2025–35.
42. Barkun AN, Almadi M, Kuipers EJ, et al. Management of nonvariceal upper gastrointestinal bleeding: Guideline recommendations from the International Consensus Group. *Ann Intern Med* 2019;171(11):805–822.
43. Gralnek IM, Stanley AJ, Morris AJ, et al. Endoscopic diagnosis and management of nonvariceal upper gastrointestinal hemorrhage (NVUGIH): European Society of Gastrointestinal Endoscopy

- (ESGE) Guideline - Update 2021. *Endoscopy*. 2021;53(3):300–332.
44. NICE Guidelines 2012. Acute Upper Gastrointestinal Bleeding in over 16s management Clinical guideline [CG141] (<https://www.nice.org.uk/guidance/cg141>).
 45. Restellini S, Kherad O, Jairath V, et al. Red blood cell transfusion is associated with increased rebleeding in patients with nonvariceal upper gastrointestinal bleeding. *Aliment Pharmacol Ther* 2013;37(3):316–322.
 46. Stokes A, Thompson C, Clegg A, et al. The influence of a simple blood transfusion policy on overtransfusion in acute upper gastrointestinal haemorrhage. *Clin Med (Lond)* 2015;15(4):325–9.
 47. Subramaniam K, Spilsbury K, Ayonrinde OT, et al. Red blood cell transfusion is associated with further bleeding and fresh-frozen plasma with mortality in nonvariceal upper gastrointestinal bleeding. *Transfusion* 2016;56(4):816–26.
 48. Chen YC, Hsiao CT, Lin LC, et al. The association between red blood cell transfusion and outcomes in patients with upper gastrointestinal bleeding. *Clin Transl Gastroenterol* 2018;9(3):138.
 49. Garcia-Tsao G, Abraldes JG, Berzigotti A, et al. Portal hypertensive bleeding in cirrhosis: Risk stratification, diagnosis, and management: 2016 practice guidance by the American Association for the study of liver diseases. *Hepatology*. 2017;65(1):310–335.
 50. European Association for the Study of the Liver. EASL Clinical Practice Guidelines for the management of patients with decompensated cirrhosis. *J Hepatol*. 2018;69(2):406–460.
 51. Liu P, Hum J, Jou J, et al. Transfusion strategies in patients with cirrhosis. *Eur J Haematol*. 2020;104(1):15–25.
 52. Boyer JL, Chatterjee C, Iber FL, et al. Effect of plasma-volume expansion on portal hypertension. *N Engl J Med*. 1966;275(14):750–755.
 53. McCormick PA, Jenkins SA, McIntyre N, et al. Why portal hypertensive varices bleed and bleed: a hypothesis. *Gut* 1995;36(1):100–103.
 54. Castaneda B, Morales J, Lionetti R, et al. Effects of blood volume restitution following a portal hypertensive-related bleeding in anesthetized cirrhotic rats. *Hepatology*. 2001;33(4):821–825.
 55. Farkas J. PulmCrit- Coagulopathy management in the bleeding cirrhotic: Seven pearls and one crazy idea. *EMCrit Blog*. <http://emcrit.org/pulmcrit/coagulopathy-bleeding-cirrhotic-inr/>. Accessed 5 June 2022.
 56. Heyes J, Kelly PA, Monaghan K, et al. A single unit transfusion policy reduces red cell transfusions in general medical in-patients. *QJM* 2017;110(11):735–739.
 57. Sung JJ, Chiu PW, Chan FKL, et al. Asia-Pacific working group consensus on non-variceal upper gastrointestinal bleeding: an update 2018. *Gut* 2018;67(10):1757–1768.
 58. Abraham NS, Barkun AN, Sauer BG, et al. American College of Gastroenterology–Canadian Association of Gastroenterology Clinical Practice Guideline: Management of Anticoagulants and Antiplatelets during Acute Gastrointestinal Bleeding and the Periendoscopic Period. *Am J Gastroenterol*. 2022;117(4):542–558.
 59. Munlemvo DM, Tobias JD, Chenault KM, et al. Prothrombin Complex Concentrates to Treat Coagulation Disturbances: An Overview with a Focus on Use in Infants and Children. *Cardiol Res*. 2022;13(1):18–26.
 60. Cohn SM, Jimenez JC, Khoury L, et al. Inability to Reverse Aspirin and Clopidogrel-induced Platelet Dysfunction with Platelet Infusion. *Cureus*. 2019;11(1):e3889.
 61. Savage SA, Sumislawski JJ, Zarzaur BL, et al. The new metric to define large-volume hemorrhage: results of a prospective study of the critical administration threshold. *J Trauma Acute Care Surg* 2015;78(2):224–9.
 62. Holcomb JB, Tilley BC, Baraniuk S, et al. Transfusion of Plasma, Platelets, and Red Blood Cells in a 1:1:1 Vs a 1:1:2 Ratio and Mortality in Patients with Severe Trauma: The Proppr Randomized Clinical Trial. *JAMA* 2015;313(5):471–82.
 63. Lau JY, Leung WK, Wu JC, et al. Omeprazole before endoscopy in patients with gastrointestinal bleeding. *N Engl J Med*. 2007;356(16):1631–40.
 64. Sreedharan A, Martin J, Leontiadis GI, et al. Proton pump inhibitor treatment initiated prior to endoscopic diagnosis in upper gastrointestinal bleeding. *Cochrane Database Syst Rev*. 2010;2010(7):CD005415.
 65. Kanno T, Yuan Y, Tse F, et al. Proton pump inhibitor treatment initiated prior to endoscopic diagnosis in upper gastrointestinal bleeding. *Cochrane Database Syst Rev*. 2022;1(1):CD005415.
 66. Bai Y, Guo JF, Li ZS. Meta-analysis: erythromycin before endoscopy for acute upper gastrointestinal bleeding. *Aliment Pharmacol Ther* 2011;34(2):166–171.
 67. Szary NM, Gupta R, Choudhary A, et al. Erythromycin prior to endoscopy in acute upper gastrointestinal bleeding: a meta-analysis. *Scand J Gastroenterol* 2011;46(7–8):920–924.
 68. Theivanayagam S, Lim RG, Cobell WJ et al. Administration of erythromycin before endoscopy in upper gastrointestinal bleeding: a metaanalysis of randomized controlled trials. *Saudi J Gastroenterol* 2013;19(5):205–210.
 69. Barkun AN, Bardou M, Martel M, et al. Prokinetics in acute upper GI bleeding: a meta-analysis. *Gastrointest Endosc* 2010;72(6):1138–1145.
 70. Rahman R, Nguyen DL, Sohail U, et al. Pre-endoscopic erythromycin administration in upper gastrointestinal bleeding: an updated metaanalysis and systematic review. *Ann Gastroenterol* 2016;29(3):312–317.
 71. Na HK, Jung HY, Seo DW, et al. Erythromycin infusion prior to endoscopy for acute nonvariceal upper gastrointestinal bleeding: A pilot randomized controlled trial. *Korean J Intern Med* 2017;32(6):1002–1009.
 72. Winstead NS, Wilcox CM. Erythromycin prior to endoscopy for acute upper gastrointestinal

- haemorrhage: a cost-effectiveness analysis. *Aliment Pharmacol Ther* 2007;26(10):1371–1377.
73. Daram S, Johnson W, Abrasley C, et al. A double blind randomized study to evaluate the use of metoclopramide before endoscopy for upper gastrointestinal bleeding. *Am J Gastroenterol* 2010;105(Suppl 1):S508.
 74. Daram SR, Garretson R. Erythromycin is preferable to metoclopramide as a prokinetic in acute upper GI bleeding. *Gastrointest Endosc* 2011;74(1):234.
 75. Estes DJ, Berera S, Deshpande AR, et al. Re-Visiting Metoclopramide to Optimize Visualization with Gastrointestinal Bleeding - Mobilizing Existing Data. *Clin Endosc.* 2019;52(5):516–517.
 76. Tripathi D, Stanley AJ, Hayes PC, et al. U.K. guidelines on the management of variceal haemorrhage in cirrhotic patients. *Gut* 2015;64(11):1680–704.
 77. de Franchis R, Baveno VI Faculty. Expanding consensus in portal hypertension: report of the Baveno VI Consensus Workshop: Stratifying risk and individualizing care for portal hypertension. *J Hepatol* 2015;63(3):743–52.
 78. Stanley AJ, Laine L. Management of acute upper gastrointestinal bleeding. *BMJ.* 2019;364:l536.
 79. Yoshiji H, Nagoshi S, Akahane T, et al. Evidence-based clinical practice guidelines for Liver Cirrhosis 2020. *J Gastroenterol.* 2021;56(7):593–619.
 80. Riha HM, Wilkinson R, Twilla J, et al. Octreotide added to a proton pump inhibitor versus a proton pump inhibitor alone in nonvariceal upper-gastrointestinal bleeds. *Ann Pharmacother* 2019;53(8):794–800.
 81. Wells M, Chande N, Adams P, et al. Meta-analysis: vasoactive medications for the management of acute variceal bleeds. *Aliment Pharmacol Ther* 2012;35(11):1267–78.
 82. Zhou X, Tripathi D, Song T, et al. Terlipressin for the treatment of acute variceal bleeding: a systematic review and meta-analysis of randomized controlled trials. *Medicine (Baltimore).* 2018;97(48):e13437.
 83. Bolognesi M, Di Pascoli M, Verardo A, et al. Splanchnic vasodilation and hyperdynamic circulatory syndrome in cirrhosis. *World J Gastroenterol.* 2014;20(10):2555–2563.
 84. Wong YJ, Tan CK, Lin Yii YL, et al. Antibiotic prophylaxis in cirrhosis patients with upper gastrointestinal bleeding: An updated systematic review and meta-analysis. *Portal Hypertension & Cirrhosis* 2022;1(3):167–177.
 85. Tandon P, Abraldes JG, Keough A, et al. Risk of bacterial infection in patients with cirrhosis and acute variceal hemorrhage, based on child Pugh class, and effects of antibiotics. *Clin Gastroenterol Hepatol.* 2015;13(6):1189–96.e2.
 86. Gao Y, Qian B, Zhang X, et al. Prophylactic antibiotics on patients with cirrhosis and upper gastrointestinal bleeding: A meta-analysis. *PLoS One.* 2022;17(12):e0279496.
 87. Bennett C, Klingenberg SL, Langholz E, et al. Tranexamic acid for upper gastrointestinal bleeding. *Cochrane Database Syst Rev* 2014;2014(11):CD006640.
 88. HALT-IT Trial Collaborators. Effects of a high-dose 24-h infusion of tranexamic acid on death and thromboembolic events in patients with acute gastrointestinal bleeding (HALT-IT): an international randomised, double-blind, placebo-controlled trial. *Lancet* 2020;395(10241):1927–1936.
 89. Aljebreen AM, Fallone CA, Barkun AN. Nasogastric aspirate predicts high-risk endoscopic lesions in patients with acute upper-GI bleeding. *Gastrointest Endosc.* 2004;59(2):172–8.
 90. Rockey DC, Ahn C, de Melo SW Jr. Randomized pragmatic trial of nasogastric tube placement in patients with upper gastrointestinal tract bleeding. *J Investig Med.* 2017;65(4):759–764.
 91. Karakonstantis S, Tzagarakis E, Kalemaki D, et al. Nasogastric aspiration/lavage in patients with gastrointestinal bleeding: a review of the evidence. *Expert Rev Gastroenterol Hepatol* 2018;12(1):63–72.
 92. Palamidessi N, Sinert R, Falzon L, et al. Nasogastric aspiration and lavage in emergency department patients with hematochezia or melena without hematemesis. *Acad Emerg Med.* 2010;17(2):126–32.
 93. Huang ES, Karsan S, Kanwal F, et al. Impact of nasogastric lavage on outcomes in acute GI bleeding. *Gastrointest Endosc* 2011;74(5):971–80.
 94. Pallin DJ, Saltzman JR. Is nasogastric tube lavage in patients with acute upper GI bleeding indicated or antiquated? *Gastrointest Endosc.* 2011;74(5):981–4.
 95. Laine L, Jensen DM. Management of patients with ulcer bleeding. *Am J Gastroenterol.* 2012;107(3):345–60;quiz 361.
 96. Gené E, Calvet X. Nasogastric intubation in patients with upper gastrointestinal bleeding? *Gastroenterol Hepatol.* 2016;39(8):497–9.
 97. Gong EJ, Hsing LC, Seo HI, et al. Selected nasogastric lavage in patients with nonvariceal upper gastrointestinal bleeding. *BMC Gastroenterol.* 2021;21(1):113.
 98. Saltzman JR. Approach to acute upper gastrointestinal bleeding in adults. In: *UpToDate, Post TW (Ed), UpToDate, Waltham, MA.* (Accessed on June 1, 2022).
 99. Dai JY, Yin L, Xu J, et al. Expert consensus on emergency diagnosis and treatment procedures for acute upper gastrointestinal bleeding. *J Acute Dis* 2020;9(6):231–43.
 100. Kim JS, Kim BW, Kim DH, et al. Guidelines for Nonvariceal Upper Gastrointestinal Bleeding. *Gut Liver.* 2020;14(5):560–570.
 101. Spiegel BM, Vakil NB, Ofman JJ. Endoscopy for acute nonvariceal upper gastrointestinal tract hemorrhage: is sooner better? A systematic review. *Arch Intern Med* 2001;161(11):1393–404.
 102. Cooper GS, Kou TD, Wong RC. Use and impact of early endoscopy in elderly patients with peptic ulcer hemorrhage: A population-based analysis. *Gastrointest Endosc* 2009;70(2):229–35.
 103. Laursen SB, Leontiadis GI, Stanley AJ, et al.

-
- Relationship between timing of endoscopy and mortality in patients with peptic ulcer bleeding: A nationwide cohort study. *Gastrointest Endosc* 2017;85(5):936–944.e3.
104. Garg SK, Anugwom C, Campbell J, et al. Early esophagogastroduodenoscopy in upper gastrointestinal bleeding: a nationwide study. *Endosc Int Open* 2017;5(5):E376–E386.
105. Xia XF, Chiu PWY, Tsoi KKF, et al. The effect of off-hours hospital admission on mortality and clinical outcomes for patients with upper gastrointestinal hemorrhage: a systematic review and meta-analysis of 20 cohorts. *United European Gastroenterol J*. 2018;6(3):367–381.
106. Siau K, Hodson J, Ingram R, et al. Time to endoscopy for acute upper gastrointestinal bleeding: results from a prospective multicentre trainee-led audit. *United European Gastroenterol J* 2019;7(2):199–209.
107. Jeong N, Kim KS, Jung YS, et al. Delayed endoscopy is associated with increased mortality in upper gastrointestinal hemorrhage. *Am J Emerg Med*. 2019;37(2):277–280.
108. Guo CLT, Wong SH, Lau LHS, et al. Timing of endoscopy for acute upper gastrointestinal bleeding: a territory-wide cohort study. *Gut*. 2021;gutjnl-2020-323054.
109. Jairath V, Kahan BC, Logan RF, et al. Outcomes following acute nonvariceal upper gastrointestinal bleeding in relation to time to endoscopy: results from a nationwide study. *Endoscopy*. 2012;44(8):723–30.
110. Lim LG, Ho KY, Chan YH, et al. Urgent endoscopy is associated with lower mortality in high-risk but not low-risk nonvariceal upper gastrointestinal bleeding. *Endoscopy* 2011;43(4):300–6.
111. Targownik LE, Murthy S, Keyvani L, et al. The role of rapid endoscopy for high-risk patients with acute nonvariceal upper gastrointestinal bleeding. *Can J Gastroenterol*. 2007;21(7):425–9.
112. Tai CM, Huang SP, Wang HP, et al. High-risk ED patients with nonvariceal upper gastrointestinal hemorrhage undergoing emergency or urgent endoscopy: a retrospective analysis. *Am J Emerg Med*. 2007;25(3):273–8.
113. Lau JYW, Yu Y, Tang RSY, et al. Timing of endoscopy for acute upper gastrointestinal bleeding. *N Engl J Med* 2020;382(14):1299–1308.
114. Kim J, Gong EJ, Seo M, et al. Timing of endoscopy in patients with upper gastrointestinal bleeding. *Sci Rep* 2022;12:6833.
115. Laursen SB, Hansen JM, Schaffalitzky de Muckadell OB. The Glasgow Blatchford score is the most accurate assessment of patients with upper gastrointestinal hemorrhage. *Clin Gastroenterol Hepatol* 2012;10(10):1130–1135.e1.
116. Tham J, Stanley A. Clinical utility of pre-endoscopy risk scores in upper gastrointestinal bleeding. *Expert Rev Gastroenterol Hepatol*. 2019;13(12):1161–1167.

KidneyEducation

Join the Mission to Fight Kidney Diseases

Explore the world's largest multilingual website created by a global team of over 100 nephrologists.

www.KidneyEducation.com

- » Read online or download the 200-page book "Save Your Kidneys" in 40 languages—completely free.
- » This comprehensive resource offers valuable information on preventing and managing common kidney problems, tailored for kidney patients and their families.
- » It's an authentic guide, prepared by nephrologists and free from any external funding.